

# Superfluid signatures in magnetar seismology

N. Andersson<sup>1</sup>, K. Glampedakis<sup>2,3</sup> & L. Samuelsson<sup>4</sup>

<sup>1</sup> *School of Mathematics, University of Southampton, Southampton SO17 1BJ, United Kingdom*

<sup>2</sup> *SISSA, via Beirut 2-4, 34014 Trieste, Italy*

<sup>3</sup> *Theoretical Astrophysics, Auf der Morgenstelle 10, University of Tuebingen, Tuebingen D-72076, Germany*

<sup>4</sup> *Nordita, Roslagstullsbacken 23, 106 91 Stockholm, Sweden*

12 December 2008

## ABSTRACT

We investigate the role of neutron star superfluidity for magnetar oscillations. Using a plane-wave analysis we estimate the effects of a neutron superfluid in the elastic crust region. We demonstrate that the superfluid imprint is likely to be more significant than the effects of the crustal magnetic field. We also consider the region immediately beneath the crust, where superfluid neutrons are thought to coexist with a type II proton superconductor. Since the magnetic field in the latter is carried by an array of fluxtubes, the dynamics of this region differs from standard magnetohydrodynamics. We show that the presence of the neutron superfluid (again) leaves a clear imprint on the oscillations of the system. Taken together, our estimates show that the superfluid components cannot be ignored in efforts to carry out “magnetar seismology”. This increases the level of complexity of the modelling problem, but also points to the possibility of using observations to probe the superfluid nature of supranuclear matter.

## 1 INTRODUCTION

Anomalous X-ray pulsars (AXPs) and Soft Gamma-ray Repeaters (SGRs) are widely believed to be magnetars; neutron stars powered by an ultrastrong magnetic field (see Woods & Thompson (2004) for a review). Observations (mainly conducted by X-ray satellites) have established basic parameters like the magnetic field intensity,  $B \sim 10^{15}$  G, and spin period,  $P \sim 10$ s, for this class of objects. They have also revealed a complex emission pattern with alternating periods of burst activity and quiescence. SGRs, which are typically more active than AXPs, are the only ones exhibiting giant flares. These flares, which are believed to be triggered by some sort of instability in the magnetic field (Duncan & Thompson 1992; Thomson & Duncan 1995), are by far the most energetic events associated with magnetars.

An exciting contribution to magnetar phenomenology was provided by the recent discovery of quasi-periodic oscillations (QPOs) in the late tail spectrum of the two giant flares (Israel et al 2005; Strohmayer & Watts 2005; Watts & Strohmayer 2006). There may also be evidence for a single QPO in the data of the third known flare, observed back in the 1970s (Barat et al 1983). The frequencies of the most prominent QPOs lie in the range 30 – 100 Hz, exactly where one would expect to find the seismic oscillation modes of the magnetar’s crust (Hansen & Cioffi 1980). This is consistent with the theoretical expectation that the energy released in a giant flare is sufficient to fracture the crust and excite its normal modes (Duncan 1998). If this interpretation of the QPOs is correct then we may have the opportunity to carry out magnetar “asteroseismology”; a comparison between theoretically predicted mode frequencies and the QPO

data, with the ultimate goal of constraining the properties of neutron star matter (Samuelsson & Andersson 2007).

Indeed, several recent papers have attempted to constrain the bulk equation of state of neutron star matter, assuming that the observed QPO frequencies are identified with the first few toroidal seismic modes of the crust (see Strohmayer (2007) for a recent review and references). This is natural as a first step, but in reality the situation is likely to be more complicated. As suggested by Levin (2006) and Glampedakis, Samuelsson & Andersson (2006) the strong magnetic field would likely couple an oscillating crust to the liquid core on a very short timescale. Then the observed QPOs would be the manifestation of the coupled crust-core dynamics rather than the dynamics of the crust alone. Possible evidence that the magnetic core plays an active role is given by the presence of a low frequency QPO in the data of the December 2004 flare in SGR 1806-20. This QPO is difficult to reconcile with the seismic mode interpretation (Israel et al 2005). It is therefore conceivable that magnetar “seismology” also probes the (much less well known) properties of the interior magnetic field. This obviously comes at a price. We now have to model global crust-core oscillations, a problem that is considerably more challenging than that of pure seismic crust modes.

Another aspect of neutron star physics, which is directly relevant to the mode interpretation of the QPOs, has received almost no attention so far. Young and mature neutron stars (older than a month or so) are sufficiently cold that the bulk of their interior liquid matter is in a superfluid state. In the crust, for densities above the nuclear drip density  $\rho \approx 4 \times 10^{11}$  g/cm<sup>3</sup>, the “dripped” neutrons are expected to form a superfluid below a temperature  $\sim 5 \times 10^9$  K. The dynamical role of these “free” neutrons could be important.

After all, they account for  $\sim 80\%$  of the total mass in the crust. Similarly, in the liquid core we expect to find both neutrons and protons in a superfluid state (below a similar threshold temperature). In the outer core, the protons most likely form a type II superconductor, provided that the interior magnetic field does not exceed a critical value  $\approx 3 \times 10^{16}$  G. As a consequence, any magnetic field that penetrates the proton plasma will form a large number of quantized magnetic fluxtubes (Baym, Pethick & Pines 1969).

It is clearly relevant to ask to what extent the physical components (the crust and the magnetic field) that play the leading role in the magnetar QPO problem are sensitive to neutron and proton superfluidity/superconductivity. The aim of this investigation is to provide an insight into this issue, and improve our understanding of the relative importance of the multifluid aspects of the magnetar oscillation problem. By carrying out a local analysis, i.e. considering uniform parameter models, we learn how the shear waves in the neutron star crust are affected by the presence of a superfluid neutron component. Similarly, a local analysis in the core tells us how the Alfvén waves are altered by the presence of the neutron superfluid (which provides the bulk of the core mass). Not surprisingly, the entrainment between neutrons and protons turns out to be the key parameter in these problems. This initial (order of magnitude) analysis serves as a useful guideline for future (more detailed) work for realistic neutron star models.

## 2 MULTIFLUID DYNAMICS OF THE CRUST

### 2.1 Lagrangian perturbation equations

We want to model linear perturbations in a neutron star crust penetrated by a superfluid neutron component. It is natural to use a Lagrangian picture to describe this problem. Hence, we combine the two-fluid Lagrangian perturbation equations (Andersson, Comer & Grosart 2004) with the relevant elastic terms and the magnetic force (Glampedakis & Andersson 2007). Since all known magnetars are slowly rotating, with periods of several seconds, it makes sense to focus on the non-rotating problem. Then we need an equation for the superfluid neutron displacement which can be written

$$(1 - \varepsilon_n) \partial_t^2 \xi_n^i + \varepsilon_n \partial_t^2 \xi_c^i + \nabla_i \delta \Phi + \xi_n^j \nabla_j \nabla_i \Phi - (\nabla_i \xi_n^j) \nabla_j \tilde{\mu}_n + \nabla_i \Delta_n \tilde{\mu}_n = 0 \quad (1)$$

We label the variables associated with the neutrons with a constituent index  $n$ .  $\Delta_n$  represents a Lagrangian variation along the neutron flow (associated with the displacement  $\xi_n^i$ ). The variable  $\varepsilon_n$  (assumed constant here) encodes the entrainment effect,  $\delta \Phi$  represents the (Eulerian) perturbation of the gravitational potential  $\Phi$  and  $\tilde{\mu}_n$  is the (specific) chemical potential for the neutrons. We have

$$\Delta_n \tilde{\mu}_n = \left( \frac{\partial \tilde{\mu}_n}{\partial n_n} \right)_{n_c} \nabla_i (n_n \xi_n^i) + \left( \frac{\partial \tilde{\mu}_n}{\partial n_c} \right)_{n_n} \nabla_i (n_c \xi_c^i) + \xi_n^i \nabla_i \tilde{\mu}_n \quad (2)$$

where  $n_n$  and  $n_c$  are the number densities of the neutrons and the baryons making up the crust nuclei, respectively. It

should be noted that we are not accounting for effects due to the presence of neutron vortices, e.g. the mutual friction and the vortex tension here. In principle, these effects will be present even in the very slowly rotating magnetars, and it will be interesting to consider them at a later stage. Our initial aim is to explore the leading order effects of this rather complicated problem.

The corresponding equation of motion for the crust nuclei can, labeling the variables associated with the nuclei with the constituent index  $c$ , be written

$$(1 - \varepsilon_c) \partial_t^2 \xi_c^i + \varepsilon_c \partial_t^2 \xi_n^i + \nabla_i \delta \Phi + \xi_c^j \nabla_j \nabla_i \Phi - (\nabla_i \xi_c^j) \nabla_j \tilde{\mu}_c + \nabla_i \Delta_c \tilde{\mu}_c = \Delta_c f_i^{\text{el}} + \Delta_c f_i^{\text{mag}} \quad (3)$$

Here  $\Delta_c$  representing the Lagrangian variation along the crust motion (associated with a displacement  $\xi_c^i$ ), and

$$\Delta_c \tilde{\mu}_c = \left( \frac{\partial \tilde{\mu}_c}{\partial n_c} \right)_{n_n} \nabla_i (n_c \xi_c^i) + \left( \frac{\partial \tilde{\mu}_c}{\partial n_n} \right)_{n_c} \nabla_i (n_n \xi_n^i) + \xi_c^i \nabla_i \tilde{\mu}_c \quad (4)$$

It is also worth noting that

$$n_n \varepsilon_n = n_c \varepsilon_c \quad (5)$$

The charged component equation includes both elastic and magnetic contributions. The former can be written

$$\Delta_c f_i^{\text{el}} = \frac{1}{\rho_c} \nabla^j \sigma_{ij} \quad (6)$$

where

$$\sigma_{ij} = \mu (\nabla_i \xi_j^c + \nabla_j \xi_i^c) - \frac{2}{3} \mu (\nabla^l \xi_l^c) \delta_{ij} \quad (7)$$

(here one should not confuse the shear modulus  $\mu$  with the chemical potentials  $\mu_x$ ). Meanwhile, the magnetic term follows from the standard electromagnetic Lorentz force. That is, in this case we have

$$f_i^{\text{mag}} = f_i^L = \frac{1}{c \rho_c} \epsilon_{ijk} J^j B^k \quad (8)$$

Eliminating the total current with the help of Ampère's law (i.e.  $J^i = (c/4\pi) \epsilon^{ijk} \nabla_j B_k$ ), this becomes

$$f_i^L = \frac{B^j}{4\pi \rho_c} (\nabla_j B_i - \nabla_i B_j) \quad (9)$$

Working out the Lagrangian variation using (Glampedakis & Andersson 2007)

$$\Delta_c \left( \frac{B^j}{\rho_c} \right) = 0 \quad (10)$$

we arrive at

$$\Delta_c B^i = -B^i \nabla_j \xi_c^j \quad (11)$$

and

$$\Delta_c B_i = B_j \nabla_i \xi_c^j - B_i (\nabla_j \xi_c^j) + B^j \nabla_j \xi_c^i \quad (12)$$

Finally, using

$$\Delta_c (\nabla_j B_i) = \nabla_j (\Delta_c B_i) - B_l \nabla_j \nabla_i \xi_c^l \quad (13)$$

we obtain from (9)

$$\Delta_c f_i^{\text{mag}} = \frac{B^j}{4\pi \rho_c} [\nabla_j (\Delta_c B_i) - \nabla_i (\Delta_c B_j)] \quad (14)$$

These are all the relations we need to solve the problem. As far as we are aware, this is the first time that the perturbation problem for combined superfluidity, elasticity and magnetic fields has been formulated. The equations we have given can be directly applied to studies of global mode oscillations of a superfluid neutron star with a crust.

For later convenience it is useful to note that we could equally well have worked with Eulerian variations. In fact, since  $f_L^i = 0$  in the background configuration we must have  $\Delta_c f_i^{\text{mag}} = \delta f_i^{\text{mag}}$ . Moreover, one can show that in the case of an incompressible fluid and a uniform background field (see below) we have

$$\Delta_c f_i^{\text{mag}} = \delta f_i^{\text{mag}} = v_A^2 \left[ (\hat{B}^j \nabla_j)^2 \xi_i^c - \hat{B}^j \hat{B}^l \nabla_l \nabla_i \xi_j^c \right] \quad (15)$$

where "hats" denote unit vectors. We have also defined the Alfvén wave velocity

$$v_A^2 = \frac{B^2}{4\pi\rho_c} \quad (16)$$

It is important to note that in the superfluid system the Alfvén velocity scales with the number density of charged nucleons, not the total baryon number density (Mendell 1998).

## 2.2 Plane-wave analysis

As a first step towards understanding the problem, let us consider the simple case of a uniform, non-rotating background. For an incompressible model, we have

$$\nabla_i \xi_x^i = 0 \longrightarrow \Delta_x \tilde{\mu}_x = 0 \quad (17)$$

Then the problem simplifies to (note that we will have  $\delta\Phi = \nabla_i \Phi = 0$  for a uniform background)

$$(1 - \varepsilon_n) \partial_t^2 \xi_i^n + \varepsilon_n \partial_t^2 \xi_i^c = 0 \quad (18)$$

$$(1 - \varepsilon_c) \partial_t^2 \xi_i^c + \varepsilon_c \partial_t^2 \xi_i^n = \Delta_c f_i^{\text{el}} + \Delta_c f_i^{\text{mag}} \quad (19)$$

We also need

$$\Delta_c f_i^{\text{el}} = v_s^2 \nabla^2 \xi_i^c \quad (20)$$

where the shear velocity,  $v_s$ , is defined by

$$v_s^2 = \frac{\mu}{\rho_c} \quad (21)$$

Note that the shear velocity scales with the number density of nucleons locked in the crust lattice, not the total nucleon number density as would be the case if there were no superfluid component. This distinction has not been made in previous work where the crust is modelled as a single component, see for example Duncan (1998); Piro (2005).

We now consider (short wavelength  $\ll$  the radius of the star) wave-propagation in this system. Making the standard plane-wave Ansatz (see Sidery, Andersson & Comer (2008) for a recent analysis of the analogous non-magnetic two-fluid problem)

$$\xi_i^x = A_i^x e^{i(\omega t + k_j x^j)} \quad (22)$$

where the index  $x$  is either  $n$  or  $c$ , we have

$$k^i A_i^x = 0 \quad (23)$$

i.e., the waves are transverse, and

$$\Delta_c f_i^{\text{mag}} = -v_A^2 k^2 \left[ (\hat{B}^j \hat{k}_j)^2 \xi_i^c - \hat{k}_i (\hat{B}_j \xi_j^c) (\hat{B}^l \hat{k}_l) \right] \quad (24)$$

From Eq. (18) we immediately get the relation

$$A_i^n = -\frac{\varepsilon_n}{1 - \varepsilon_n} A_i^c \quad (25)$$

Using this in Eq. (19) we arrive at

$$\left[ \frac{\omega^2}{\varepsilon_*} - v_s^2 k^2 - v_A^2 k^2 (\hat{B}_j \hat{k}^j)^2 \right] A_i^c = -v_A^2 k^2 (\hat{B}_j \hat{k}^j) (\hat{B}^l A_l^c) \hat{k}_i \quad (26)$$

where we have introduced

$$\varepsilon_* = \frac{1 - \varepsilon_n}{1 - \varepsilon_n - \varepsilon_c} \quad (27)$$

Defining

$$\omega_0^2 = v_s^2 k^2 \quad (28)$$

the frequency of "pure" elastic waves, and the Alfvén-wave frequency

$$\omega_A^2 = v_A^2 k^2 \quad (29)$$

we have an equation for  $A_i^c$ ,

$$\left[ \frac{\omega^2}{\varepsilon_*} - \omega_0^2 - (\hat{B}_j \hat{k}^j)^2 \omega_A^2 \right] A_i^c = -\omega_A^2 (\hat{B}_j \hat{k}^j) (\hat{B}^l A_l^c) \hat{k}_i \quad (30)$$

In order to arrive at the final dispersion relation we first note that contracting the above equation with  $\hat{k}^i$  leads to the constraint

$$\omega_A^2 (\hat{B}_j \hat{k}^j) (\hat{B}^l A_l^c) = 0 \quad (31)$$

Thus, we can either choose to look for solutions where  $k^i$  is orthogonal to the local magnetic field or we see that the polarisation  $A_i^c$ , and hence  $A_i^n$ , must be orthogonal to both  $k^i$  and  $B^i$ . Since the right hand side of (30) vanishes in both cases we find that all non-trivial solutions must be such that

$$\frac{\omega^2}{\varepsilon_*} - \omega_0^2 - (\hat{B}_j \hat{k}^j)^2 \omega_A^2 = 0 \quad (32)$$

That is, we have the general dispersion relation

$$\omega^2 = \omega_0^2 \varepsilon_* \left[ 1 + (\hat{B}_j \hat{k}^j)^2 \frac{\omega_A^2}{\omega_0^2} \right] \quad (33)$$

Note that, in the degenerate case when  $B_i k^i = 0$  we cannot uniquely determine the polarisation; it can lie in any direction in the plane orthogonal to  $k^i$ . Also, it is clear that such waves do not depend on the magnetic field at all. Generically the polarisation is, however, well defined (up to scale since we have a homogeneous system) to be orthogonal to both  $k^i$  and  $B^i$ .

We need to estimate the magnitude of the various terms. Let us first focus on the entrainment. Using (5) we find that

$$1 - \varepsilon_n - \varepsilon_c = 1 - \frac{\varepsilon_n}{x_c} \quad (34)$$

where  $x_c = \rho_c/\rho$ . We can express this in terms of the effective mass of the free neutrons,  $m_n^*$ . Then we have (see Prix, Comer & Andersson (2002) for a discussion of the analogous problem in a superfluid neutron star core)

$$\varepsilon_n = 1 - \frac{m_n^*}{m_n} \quad (35)$$

and it follows that

$$\varepsilon_\star = x_c \left(1 - x_n \frac{m_n}{m_n^*}\right)^{-1} = \frac{x_c}{\chi} \quad (36)$$

It is easy to show that  $\chi^{-1/2}$  encodes the difference between the superfluid result and the standard result for a single component crust, i.e. with  $\rho_c \rightarrow \rho$  in (21). This may be the most meaningful comparison to make, since all previous studies of crust oscillations have assumed the single component model.

What do we learn from these results? First of all, we see that in the limit  $m_n^* \rightarrow m_n$ , when the medium effects that lead to the effective mass differing from the bare mass are not so great, we have  $\varepsilon_\star \rightarrow 1$  and  $\chi \rightarrow x_c$ . The waves in such a system, cf. (33), are the usual shear waves with a (as we will see later) relatively small magnetic correction. Of course, the results could still differ significantly from the standard single component model. The largest effect that one would expect would be, for  $x_c \approx 0.8$  cf. Figure 1, a frequency increase by about a factor of 2. However, the effective mass is expected to be larger than the bare mass so let us consider the opposite limit, which may well apply in parts of the neutron star crust [see, for example, Chamel (2006)]. Then we have  $m_n^* \gg m_n$ . Using also  $x_c \leq 1$ , we see that  $\varepsilon_\star \approx x_c$  or  $\chi \rightarrow 1$ . In this limit, it is very difficult for the free neutrons to move relative to the crust. The upshot of this is that the waves in the system tend to the frequency predicted for a pure elastic crust without a superfluid component.

These two extremes show that the presence of the superfluid in the neutron star crust can have a significant effect on the frequencies of waves in the system. According to the data in Figure 1, the combined effect is at the 10% level (compared to the single component crust result). The results clearly show that the superfluid component must be considered if we want to develop high precision magnetar crust seismology. Of course, in reality we are mainly interested in the global oscillations. Then the local effects that we have worked out will be (in some sense) averaged throughout the crust. One may expect this to decrease the role of the superfluid since the effective neutron mass may only be large in parts of the crust. Of course, the real answer requires a detailed mode calculation. This problem remains to be solved. In order to provide reliable results, such an effort should draw on more complete studies of the entrainment for the crust superfluid. One should also worry about the relevance of vortex pinning and the mutual friction.


Finally, let us discuss the relative importance of the magnetic field. Scaling to "typical" values, we have

$$\mu \approx 10^{13} \left(\frac{v_s}{10^8 \text{ cm/s}}\right)^2 \left(\frac{\rho}{10^{14} \text{ g/cm}^3}\right) \text{ dyne/cm}^2 \quad (37)$$

Then, it follows from (33) that we need to consider

$$\left(\frac{\omega_A}{\omega_0}\right)^2 \approx 0.08 \left(\frac{v_s}{10^8 \text{ cm/s}}\right)^{-2} \left(\frac{\rho}{10^{14} \text{ g/cm}^3}\right)^{-1} \left(\frac{B}{10^{15} \text{ G}}\right)^2 \quad (38)$$

This shows that we can safely ignore the magnetic effects in the high-density region of the crust, cf. Figure 1. In order for the magnetic term to dominate at the base of the crust we need  $B \sim 10^{16}$  G, stronger than the field strength inferred for magnetars. Of course, one has to be a little bit careful here. First of all, it is entirely possible that the interior field



**Figure 1.** This figure illustrates the density dependence of the different parameters that affect the wave propagation in the crust. We show, as functions of the total baryon number density  $n_b$ , the superfluid neutron fraction  $x_n = n_n/n_b$  (dash-dot, left scale) for the equation of state discussed by Douchin & Haensel (2001), a fit to the effective neutron mass  $m_n^*/m_n$  (solid, right scale) based on the numerical results of Chamel (2006) (data points indicated by diamonds), and the ratio between the Alfvén and the shear wave speeds  $(v_A/v_s)^2$  (dashed, left scale). The overall effect that the presence of the crust superfluid has on the local wave propagation, compared to the standard single component crust, is (as discussed in the main text) represented by  $\chi^{-1/2}$  (solid, left scale). The horizontal dashed grey line indicates unity on the left scale.

is stronger than the exterior dipole field which leads to the observed braking of the magnetar spin. Secondly, (38) indicates that the magnetic terms will dominate as we approach the surface of the star. However, our analysis breaks down completely in the surface region. Basically, the MHD approximation is only valid as long as the Alfvén wave speed is significantly below the speed of light. In order for this to be the case, we require

$$\rho \gg 10^8 \left(\frac{B}{10^{15} \text{ G}}\right)^2 \text{ g/cm}^3 \quad (39)$$

When this is no longer true, as it will not be in a transition region near the surface of any neutron star, one must consider the complete Maxwell equations. It is interesting we compare (39) to the estimated density for the top of the crust, cf. for example Eq. (1) from Piro (2005),

$$\rho_{\text{top}} \approx 2.3 \times 10^9 \left(\frac{T}{3 \times 10^8 \text{ K}}\right)^3 \left(\frac{26}{Z}\right)^6 \left(\frac{A}{56}\right) \text{ g/cm}^3 \quad (40)$$

where  $A$  is the number of nucleons and  $Z$  is the charge per ion, respectively. This density is scaled to iron, which means that the fiducial values should be relevant near the top of the crust. This suggests that MHD remains a valid approximation throughout a magnetar crust. The transition to electromagnetism takes place in the neutron stars envelope. The nature of this transition is an important problem that deserves more attention.

### 3 MULTIFLUID DYNAMICS OF THE CORE

#### 3.1 Lagrangian perturbation equations

The magnetohydrodynamics in the core is, in the simplest case, formulated in terms of three distinct fluids associated with the neutrons, protons and electrons. The former two particle species are expected to be superfluid and superconducting, respectively. Due to the smallness of the electron mass, the electron fluid degree of freedom can be suppressed and the MHD equations effectively lead to a two-fluid model (Mendell 1998). The protons in the outer core are expected to form a type II superconductor (Baym, Pethick & Pines 1969) which means that the magnetic field is carried by a large number of fluxtubes, each with a flux quantum  $\phi_0 = hc/2e$ . This should be the case provided the magnetic field is below a critical value  $H_{c2} \approx 3 \times 10^{16}$  G (Baym, Pethick & Pines 1969). This critical value represents the field strength at which the magnetic fluxtubes overlap and can no longer be treated individually. Above this threshold the magnetic field behaves "classically". It should be noted that, even though the critical field is large, this is not an unrealistic possibility for magnetars given that the magnetic field in the interior could be considerably higher than the exterior dipole field. Hence, one ought to consider both superconducting and normal protons. The latter case is, however, trivial. The desired result follows immediately from the previous section, e.g. (33), if we take the limit  $\mu \rightarrow 0$ .

For a non-rotating star, we again neglect the vortex-mediated mutual friction and the neutron vortex tension. Omitting also a small entrainment induced magnetic term that originates from the London field by means of which the proton superconductor rotates [see Glampedakis, Samuelsson & Andersson (2008) for discussion], the superfluid neutron dynamics is still governed by (1).

The combined proton-electron dynamics is a little bit more complicated. As discussed by Glampedakis, Samuelsson & Andersson (2008) the relevant equation of motion takes the form

$$\begin{aligned} (\partial_t + v_c^j \nabla_j) (v_i^c + \varepsilon_c w_i^{nc}) + \nabla_i (\tilde{\mu} + \Phi) + \varepsilon_c w_j^{nc} \nabla_i v_c^j \\ = \frac{1}{\rho_c} (f_i^L + t_i^c) \end{aligned} \quad (41)$$

where  $v_c^i$  and  $v_n^i$  are the velocities and  $w_n^i = v_n^i - v_c^i$ , and we have (again) neglected the mutual friction. There are two force terms on the right-hand side of this equation. The first,  $f_i^L$ , is the usual electromagnetic Lorentz force given by (8). The second,  $t_i^c$ , represents the smooth-averaged tension of the magnetic fluxtubes. Remarkably, the Lorentz force does not play a role in the final superconducting MHD equations. As discussed by Glampedakis, Samuelsson & Andersson (2008) [see also (Mendell 1998)] it is exactly cancelled by a term originating from the fluxtube tension. In the case of a non-rotating star this leads to the magnetic force taking the form

$$f_i^L + t_i^c = \frac{m_c c}{4\pi e} \mathcal{W}_c^j \left[ \nabla_j (H_{c1} \hat{\mathcal{W}}_i^c) - \nabla_i (H_{c1} \hat{\mathcal{W}}_j^c) \right] \quad (42)$$

where the lower critical magnetic field  $H_{c1} = H_{c1}(\rho_c) \approx 10^{15}$  G.

We have defined the vector  $\mathcal{W}_c^i$ , representing the (av-

eraged) canonical proton vorticity (Prix 2005). This means that we have<sup>1</sup>

$$\mathcal{W}_c^i \approx \frac{e}{m_c c} B^i \quad (43)$$

For a uniform incompressible model, this means that

$$\delta f_i^{\text{mag}} = \frac{1}{\rho_c} \delta (f_i^L + t_i^c) \approx \frac{H_{c1}}{4\pi \rho_c} B^j \left[ \nabla_j \delta \hat{B}_i - \nabla_i \delta \hat{B}_j \right] \quad (44)$$

In order to close the system of the MHD equations we need to provide a relation between the magnetic field and the fluid velocity. This relation follows from the magnetic induction equation. Neglecting the coupling forces between the electrons and the neutron and proton fluids (see Glampedakis, Samuelsson & Andersson (2008) for discussion) the induction equation takes the standard form,

$$\partial_t B^i \approx \epsilon^{ijk} \epsilon_{klm} \nabla_j (v_c^l B^m) \quad (45)$$

Its perturbed form is given by (10) from which we obtain the Eulerian perturbation of the magnetic field,

$$\delta B^i = B^j \nabla_j \xi_c^i - \nabla_j (\xi_c^j B^i) \quad (46)$$

Using this result, we find (for a uniform background and incompressible perturbations)

$$\delta f_{\text{mag}}^i \approx c_A^2 \hat{B}^j \hat{B}^l (g^{ik} - \hat{B}^i \hat{B}^k) \nabla_j \nabla_l \xi_c^k \quad (47)$$

where

$$c_A^2 = \frac{H_{c1} B}{4\pi \rho_c} \quad (48)$$

We now have all the relations we need to discuss short wavelength waves in the superfluid/superconducting system.

#### 3.2 Plane-wave analysis

Most of the analysis works out exactly as in the crust problem (obviously in the  $\mu \rightarrow 0$  limit). The only difference is the form of the perturbed magnetic force. With the plane-wave assumption, we see that

$$\Delta_c f_i^{\text{mag}} = -c_A^2 k^2 (\hat{B}^j \hat{k}_j)^2 \left[ \xi_i^c - (\hat{B}^l \xi_l^c) \hat{B}_i \right] \quad (49)$$

Comparing the magnetic forces in the normal and superconducting cases (equations (24) and (49), respectively) we note some important differences. The characteristic speeds are obviously different,  $c_A^2 = (H_{c1}/B) v_n^2$ . This is a well-known effect (Easson & Pethick 1977; Mendell 1998). The two velocities would differ by a factor  $\sim 10^3$  for a canonical pulsar with  $B = 10^{12}$  G. However, the effect will not be as dramatic for magnetars for which  $B \approx H_{c1} \approx 10^{15}$  G. Another difference arises from the second term in (49). This term is present provided  $\hat{B}^l \xi_l^c \neq 0$ , or in other words when the wave-vector  $k^i$  does not point along the direction of the magnetic field.

<sup>1</sup> Strictly speaking, this expression is only valid for the non-rotating background. Perturbations of  $\mathcal{W}_c^i$  will also contain the perturbed fluid velocities. However, as long as we are focussing on the leading order contributions these can be neglected. That this is a legitimate approximation is easy to see since the characteristic frequency

$$\frac{e}{m_c c} B \approx 10^{19} \left( \frac{B}{10^{15} \text{ G}} \right) \text{ s}^{-1}$$

is much higher than any other relevant frequency in the problem.

As a result, the magnetic force for superconducting protons has a non-zero component along the magnetic field. Meanwhile, we have  $B_i \Delta_c f_{\text{mag}}^i = 0$  in the normal proton case, cf. (24). In the plane-wave problem the terms that give rise to this difference vanish identically. However, this rather subtle difference in the magnetic forces will be relevant in many less idealised situations, e.g. for global mode oscillations. Hence, one must be careful before using intuition gained from standard MHD problems in the case of a superconducting core. There is certainly more<sup>2</sup> to the problem than a simple "replacement"  $B^2 \rightarrow BH_{c1}$ .

Combining the perturbation equations as in the previous section, we readily arrive at

$$\left[ \frac{\omega^2}{\varepsilon_*} - c_A^2 k^2 (\hat{k}_j \hat{B}^j)^2 \right] A_c^i = -c_A^2 k^2 (\hat{k}_j \hat{B}^j)^2 (\hat{B}_j A_c^j) \hat{B}^i \quad (50)$$

Working things out as in the crust case, we project this equation onto  $\hat{k}_i$ . Since  $k_i A_c^i = 0$  it then follows that we have the constraint

$$c_A^2 (\hat{k}_j \hat{B}^j)^3 (\hat{B}_j A_c^j) = 0 \quad (51)$$

This has the same implications as in the crust problem. It immediately follows that the right-hand side of (50) must vanish. Hence, we have the dispersion relation,

$$\omega^2 = \varepsilon_* c_A^2 k^2 (\hat{k}_j \hat{B}^j)^2 = \varepsilon_* \frac{H_{c1} B}{4\pi\rho_c} k^2 (\hat{k}_j \hat{B}^j)^2 \quad (52)$$

Our main interest here concerns the role of the superfluid neutron component. Its presence is reflected by the entrainment factor in (52). To quantify its relevance, we express the entrainment in terms of the effective proton mass, i.e. we use  $\varepsilon_c = 1 - m_p^*/m_p$ . Then it follows, since the proton fraction in the core is small, that

$$\omega^2 \approx \frac{m_p}{m_p^*} c_A^2 k^2 \quad (53)$$

Since it is expected that  $0.3 < m_p^*/m_p < 0.7$  (see Prix, Comer & Andersson (2002) for discussion) we see that the presence of the superfluid neutrons will lead to a  $\sim 20 - 80\%$  increase in the frequency of the core waves. This effect is large enough that cannot be neglected. It may, in fact, also be observable. If one accepts the argument that the magnetic field couples motion in the crust to the core, and that the core fluid is therefore partaking on the oscillation, then the entrainment will affect the observed frequencies.

We cannot at this point say much about the global oscillations of a magnetic neutron star core; it is a problem that remains to be solved in detail. It is complicated by the likely presence of an "Alfvén continuum" (Levin 2007). At this point it is not clear to what extent the continuum prevails in more detailed neutron star models. However, it is easy to see how the presence of the superfluid will manifest itself in the continuum toy-model considered by Levin (2007). The frequency range of the continuum will simply scale according to (52).

<sup>2</sup> For an interesting recent discussion on how superconductivity may affect the stability properties of the star, see Akgün & Wasserman (2008).

## 4 CONCLUDING REMARKS

In this paper we have investigated the role of neutron star superfluidity for magnetar oscillations. The results impact on attempts to use data from observed quasiperiodic oscillations in the tails of magnetar flares to place constraints on neutron star parameters, see for example Samuelsson & Andersson (2007). Using a plane-wave analysis we estimated the effects of the neutron superfluid in the elastic crust region. This, the first ever, analysis of the combined magnetic-elastic-superfluid crust problem demonstrated that the superfluid imprint is likely to be more significant than the effects of the crustal magnetic field. This is, of course, assuming that the SGR flare mechanism does not deposit sufficient heat in the crust to raise the system above the superfluid transition temperature. Available estimates, e.g. Kouveliotou et al (2003), suggest that this is unlikely. We also considered the region immediately beneath the crust, where superfluid neutrons are thought to coexist with a type II proton superconductor. Since the magnetic field in the latter is carried by an array of fluxtubes, see Glampedakis, Samuelsson & Andersson (2008) for a recent discussion, the dynamics of this region differs from standard magnetohydrodynamics. We showed that the presence of the neutron superfluid (again) affects the oscillations of the system. This accords well with previous results of, in particular, Mendell (1998).

Our estimates show that the superfluid components cannot be ignored in efforts to carry out magnetar seismology. This increases the level of complexity of the modelling problem, but also points to the exciting possibility of using observations to probe the superfluid nature of supranuclear matter. Future work needs to extend our analysis to consider the global oscillations of magnetic-superfluid-elastic neutron stars. This is a very interesting problem because, in addition to enabling a more detailed seismology analysis, it may also provide insight into rotational glitches in magnetars (Dib, Kaspi & Gavril 1992). It is generally believed that superfluidity plays a key role in radio pulsar glitches. Recent developments in modelling these events is showing some promise (Glampedakis & Andersson 2008), and it would obviously be highly relevant to extend this analysis to strongly magnetised systems.

## ACKNOWLEDGEMENTS

We would like to thank Anna Watts for helpful discussions. This work was supported by STFC in the UK through grant number PP/E001025/1. KG is supported by the German Science Foundation (DFG) via SFB/TR7.

## REFERENCES

- Akgün, T., Wasserman, I., 2008, MNRAS, 383, 1551
- Andersson N., Comer, G.L, Grosart, K., 2004, MNRAS 355, 918
- Baym, G., Pethick, C., Pines, D., 1969, Nature **224**, 673
- Barat C. et al, 1983, A & A, 126, 400
- Chamel, N., 2006, Nucl. Phys. A, 773, 263
- Douchin F., Haensel P., 2001, Astron. Astrophys., 380, 151
- Dib, R., Kaspi V.M., Gavril F.T., 2008, ApJ, 673, 1044

- Duncan R.C. and Thompson C., 1992, ApJ, 392, L9  
Duncan R.C., 1998, ApJ, 498, L45  
Eassons, I., Pethick, C.J., 1977, Phys. Rev. D 16, 275  
Glampedakis, K., Andersson, N., 2007, MNRAS 377, 630  
Glampedakis, K., Andersson, N., 2008, preprint  
arXiv:0806.3664  
Glampedakis, K., Samuelsson, L., Andersson, N., 2006,  
MNRAS 371, L74  
Glampedakis, K., Samuelsson, L., Andersson, N., 2008, in  
preparation  
Hansen C.J. and Cioffi D.F., 1980, ApJ, 238, 740  
Israel G.L. et al, 2005, ApJ, 628, L53  
Kouveliotou C., et al, 2003, ApJ 596, L79  
Levin Y., 2006, MNRAS, 368, L35  
Levin Y., 2007, MNRAS, 377, 159  
Mendell, G., 1998, MNRAS 296, 903  
Piro A.L., 2005, Ap. J. Lett, 634, 153  
Prix, R., Comer, G.L., Andersson, N., 2002, Astron. As-  
trop. 381, 178  
Prix, R., 2005, Phys. Rev. D, 71, 083006  
Samuelsson, L., Andersson, N., 2007, MNRAS, 374, 256  
Sidery, T., Andersson, N., Comer, G.L., 2008, MNRAS 385,  
335  
Strohmayer T.E., and Watts A.L., 2005, ApJ, 632, L111  
Strohmayer T.E., 2007, in “Astrophysics of Com-  
pact Objects” proceedings, Huangshan, China (preprint  
arXiv:0710.2475)  
Thomson C., and Duncan R.C., 1995, MNRAS, 275, 255  
Watts A.L., and Strohmayer T.E., 2006, ApJ, 637, L117  
Woods P.M., and Thompson C., 2004, in “Compact Stellar  
X-ray Sources” eds. W.H.G. Lewin and M. Van der Klis  
(preprint astro-ph/0406133)